

**ST. CLAIR CATHOLIC DISTRICT SCHOOL BOARD
POLICIES AND PROCEDURES
SECTION C: STUDENTS**

Appendix D

**PREVALENT MEDICAL CONDITION — ANAPHYLAXIS
Plan of Care**

STUDENT INFORMATION

Student Name _____	Date Of Birth _____
Ontario Ed. # _____	Age _____
Grade _____	Teacher(s) _____

EMERGENCY CONTACTS (LIST IN PRIORITY)

NAME	RELATIONSHIP	DAYTIME PHONE	ALTERNATE PHONE
1.			
2.			
3.			

KNOWN LIFE-THREATENING TRIGGERS

CHECK (✓) THE APPROPRIATE BOXES

Food(s): _____ Insect Stings: _____
 Other: _____
 Epinephrine Auto-Injector(s) Expiry Date (s): _____
 Dosage: EpiPen® Jr. 0.15 mg EpiPen® 0.30 mg Location Of Auto-Injector(s): _____

 Previous anaphylactic reaction: **Student is at greater risk.**
 Has asthma. **Student is at greater risk.** If student is having a reaction and has difficulty breathing, give epinephrine before asthma medication.
 Any other medical condition or allergy?

**ST. CLAIR CATHOLIC DISTRICT SCHOOL BOARD
POLICIES AND PROCEDURES
SECTION C: STUDENTS**

DAILY/ROUTINE ANAPHYLAXIS MANAGEMENT

SYMPTOMS

A STUDENT HAVING AN ANAPHYLACTIC REACTION MIGHT HAVE ANY OF THESE SIGNS AND SYMPTOMS:

- **Skin system:** hives, swelling (face, lips, tongue), itching, warmth, redness.
- **Respiratory system** (breathing): coughing, wheezing, shortness of breath, chest pain or tightness, throat tightness, hoarse voice, nasal congestion or hay fever-like symptoms (runny, itchy nose and watery eyes, sneezing), trouble swallowing.
- **Gastrointestinal system** (stomach): nausea, vomiting, diarrhea, pain or cramps.
- **Cardiovascular system** (heart): paler than normal skin colour/blue colour, weak pulse, passing out, dizziness or lightheadedness, shock.
- **Other:** anxiety, sense of doom (the feeling that something bad is about to happen), headache, uterine cramps, metallic taste.

EARLY RECOGNITION OF SYMPTOMS AND IMMEDIATE TREATMENT COULD SAVE A PERSON'S LIFE.

Avoidance of an allergen is the main way to prevent an allergic reaction.

Food Allergen(s): eating even a small amount of a certain food can cause a severe allergic reaction.

Food(s) to be avoided: _____

Safety measures: _____

Insect Stings: (Risk of insect stings is higher in warmer months. Avoid areas where stinging insects nest or congregate. Destroy or remove nests, cover or move trash cans, keep food indoors.)

Designated eating area inside school building

Safety measures:

Other information:

**ST. CLAIR CATHOLIC DISTRICT SCHOOL BOARD
POLICIES AND PROCEDURES
SECTION C: STUDENTS**

EMERGENCY PROCEDURES (DEALING WITH AN ANAPHYLACTIC REACTION)

ACT QUICKLY. THE FIRST SIGNS OF A REACTION CAN BE MILD, BUT SYMPTOMS CAN GET WORSE QUICKLY.

STEPS

1. Give epinephrine auto-injector (e.g. EpiPen®) at the first sign of known or suspected anaphylactic reaction.

2. Call 9-1-1 or local emergency medical services. Tell them someone is having a life-threatening allergic reaction.

3. Give a second dose of epinephrine as early as five (5) minutes after the first dose if there is no improvement in symptoms.

4. Go to the nearest hospital immediately (ideally by ambulance), even if symptoms are mild or have stopped. The reaction could worsen or come back, even after treatment. Stay in the hospital for an appropriate period of observation as decided by the emergency department physician (generally about 4 — 6 hours).

5. Call emergency contact person; e.g. Parent(s)/Guardian(s).

**ST. CLAIR CATHOLIC DISTRICT SCHOOL BOARD
POLICIES AND PROCEDURES
SECTION C: STUDENTS**

HEALTHCARE PROVIDER INFORMATION (OPTIONAL)

Healthcare provider may include: Physician, Nurse Practitioner, Registered Nurse, Pharmacist, Respiratory Therapist, Certified Respiratory Educator, or Certified Asthma Educator.

Healthcare Provider's Name: _____

Profession/Role: _____

Signature: _____ Date: _____

Special Instructions/Notes/Prescription Labels:

If medication is prescribed, please include dosage, frequency and method of administration, dates for which the authorization to administer applies, and possible side effects.

*This information may remain on file if there are no changes to the student's medical condition.

AUTHORIZATION/PLAN REVIEW

INDIVIDUALS WITH WHOM THIS PLAN OF CARE IS TO BE SHARED

1. _____ 2. _____ 3. _____

4. _____ 5. _____ 6. _____

Other Individuals To Be Contacted Regarding Plan Of Care:

Before-School Program Yes No _____

After-School Program Yes No _____

School Bus Driver/Route # (If Applicable) _____

Other: _____

This plan remains in effect for the 20__ — 20__ school year without change and will be reviewed on or before: _____. (It is the parent(s)/guardian(s) responsibility to notify the principal if there is a need to change the plan of care during the school year).

Parent(s)/Guardian(s): _____ Date: _____
Signature

Student: _____ Date: _____
Signature

Principal: _____ Date: _____
Signature

ST. CLAIR CATHOLIC DISTRICT SCHOOL BOARD
POLICIES AND PROCEDURES
SECTION C: STUDENTS

Using **EpiPen[®]** / **EpiPen[®] Jr**
is as easy as 1-2-3

1. Remove yellow or green cap from carrying case
 - Grasp unit with black tip pointing downward
 - Pull off grey activator cap



2. Jab black tip firmly into outer thigh so it “clicks” AND HOLD on thigh approximately 10 seconds
 - Massage injected area for 10 seconds



3. Seek medical attention



After Using **EpiPen[®]** / **EpiPen[®] Jr**
Follow 3 Easy Safety Steps:

1. Carefully place used auto-injector, needle-end first, into storage tube
2. Screw cap of carrying case on completely.
 - This automatically bends needle back and secures pen so it won't fall out of tube
3. Take unit with you to hospital Emergency Department

